RECONNAISSANCE MAP AND DESCRIPTION OF THE CHETASLINA VOLCANIC DEBRIS FLOW (NEW NAME). SOUTHEASTERN COPPER RIVER BASIN AND ADJACENT AREAS, SOUTH-CENTRAL ALASKA

Lynn A. Yehle and Donald R. Nichols

EXPLANATION

X? (1) ISOLATED EXPOSURE OF CHETASLINA VOLCANIC DEBRIS FLOW--Queried where thought to be present. Numbers refer to analyzed samples (see fig. 1 and table 1)

▲ K-AR-DATED ANDESITE SAMPLES--Age no older than about 200,000 years at the 95-percent confidence level (J. D. Obradovich, written commun. 1979; U.S. Geological Survey samples 3763 and 3810)

AREA OF CLOSELY SPACED EXPOSURES OF CHETASLINA VOLCANIC DEBRIS FLOW--Ruled where concealed or inferred

LAVA FLOWS--Probably overwhelmingly andesite. Wide ruled where inferred POSSIBLE PRINCIPAL ROUTE OF CHETASLINA VOLCANIC DEBRIS FLOW ON UPPER PART

POSSIBLE PRINCIPAL ROUTE OF LAVA FLOWS THAT PRESUMABLY OVERLIE THE CHETASLINA VOLCANIC DEBRIS FLOW

OF MOUNT WRANGELL

----- MARGIN OF ICEFIELDS AND LARGE GLACIERS · ABANDONED CHANNEL OF ANCESTRAL COPPER

DISCUSSION Introduction

The volcanoes comprising most of the glacier-clad Wrangell Mountains form a prominent part of the landscape of south-central Alaska. The volcanoes in the western part of the Wrangell Mountains flank the Copper River basin and Chitina River valley and have been the source of widespread outpourings of lava, volcaniclastic materials, and tephra. Debrisflow deposits probably ranging in type from avalanches to mudflows (lahars) are among the numerous deposits of volcaniclastic materials. The term, debris flow, is used in the sense of Sharp and Nobles (1953) to designate any rapid flowage involving debris of various kinds and conditions.

One volcanic debris-flow deposit, thought to originate from Mount Wrangell, is largely buried and discontinuous but is clearly unique, because it contains multihued clasts, some of very large size; many clasts seem to be derived from hydrothermally altered volcanic rocks. The deposit, part of which was described earlier as volcanic rubble (Nichols and Yehle, 1969), is here-named informally the Chetaslina volcanic debris flow after excellent, although discontinuous, exposures in bluffs along the middle reaches of the Chetaslina and East Fork Chetaslina Rivers. The debris flow was observed at numerous other localities in the map area during the course of reconnaissance geologic mapping by us and others (Schrader and Spencer, 1901; D. R. Nichols, written commun., 1954-58; L. A. Yehle, written commun., 1960-62, 1966: Nichols and Yehle, 1969; Miller and Smith, 1976; K. M. Johnson, oral commun., 1979; T. P. Miller, R. L. Smith, D. H. Richter, M. A. Lanphere, and

G. B. Dalrymple, written commun., 1979; Yehle The only other extensive volcanic debrisflow deposit described in the map area is the Sanford volcanic mudflow named for exposures along the Sanford River (Ferrians and others, 1958; Ferrians and Nichols, 1965, p. 104). The Sanford mudflow is well exposed along the Copper River in bluffs which are 45 km upstream from the nearest known deposits of the Chetaslina volcanic debris flow.

Other volcanic debris-flow exposures in the map area include the volcanic avalanche deposits of Nadina valley exposed southwest of Mount Drum (Richter and others, 1979), deposits exposed along the middle reach of the Nadina River (D. R. Nichols, written commun., 1956), along the Copper River near the mouths of the Nadina and Dadina Rivers (D. R. Nichols, written communs., 1956, 1958), and about 7 km northeas of the mouth of the Klawasi River (D. R. Nichols, written commun., 1955). The relationship of these deposits to the Chetaslina volcanic debris flow is not known. Some multicolored cobbles and boulders of volcanic rocks were observed in the upper part of a mid-bluff diamicton deposit, apparently of glacial origin, along the Chetaslina River. The deposit is interpreted to be a local influx of different volcanic material from Mount Wrangell upon glaciers within the Chetaslina River drainage basin. The event probably occurred a substantial time before the emplacement of the Chetaslina volcanic debris flow as evidenced by the large stratigraphic separation between the diamicton and the debris flow. We acknowledge the help of U.S. Geological Survey colleagues H. C. Starkey and Fred Hildebrand for X-ray analyses, R. L. Smith fo petrographic analyses, and R. H. Bennett and

Geographic setting of Mount Wrangell and the area of deposition

Eric Smirnow for mechanical size analyses of

unconsolidated deposits.

The volcanoes of the Wrangell Mountains, several of which rise to altitudes of over 4900 m, are mostly andesitic, and probably began developing in Miocene time (Deininger, 1972, p. 19). Among the volcanoes, only Mount Wrangell, one of the world's largest andesite volcanoes (4317 m high) (Furst, 1968, p. 13), is known to exhibit any volcanic activity. In 1974, North Crater on the north part of the top of Mount Wrangell, warmed up considerably and

Geographic name submitted to the U.S. Board of Geographic Names for approval in 1979.

melted large quantities of nearby ice (Benson and Motyka, 1976, 1977, 1979). North and Wrangell Craters are the most prominent examples of modern-day activity and form part of a discontinuous group of small craters that extend around part of the rim of the glacier-filled Mount Wrangell caldera. The caldera about 16.5 km² in area is partly breached on its south side, and tributaries of Long Glacier are enlarging the breach. On the western flank of the upper part of the volcano, there are several small fumaroles (Mendenhall, 1903, p. 400, 403; Dunn, 1909, p. 501, 503; Furst, 1968, p. 3; D. R. Nichols, written commun., 1958). Terrain flanking much of the western and southwestern part of the Wrangell Mountains generally slopes downward away from the mountains and toward the Copper River basin and its southeastward-trending extension, the Chitina River valley. Major streams have cut as deeply as about 200 m into the floor of the basin and have exposed a complex of geologic materials, largely

unconsolidated, that includes remnants of the

commonly is pinkish tan to pink to pinkish gray

and is composed of variably sized fragments of

volcanic material ranging from pebbly sand to

silt. Size analyses of eight representative

samples (fig. 1) from six widely separated

localities indicate the median size of the

matrix is 0.8 phi (within coarse sand size),

the twenty-fifth quartile is -2.3 phi (within

seventy-fifth quartile is 3.6 phi (with very

fine sand size). None of the size analyses were

from material that had been obviously hydrother-

mally altered; three samples (Nos. 1, 2A, and

of the Chetaslina debris flow has not been

2B) were from deposits which resembled diamicton

of possible glacial origin. Although the matrix

studied mineralogically on a systematic basis, a

few samples of matrix material that appeared to

be hydrothermally altered were analyzed by X-ray

diffraction techniques (table 1). A distinctive

aspect of the mineralogical analyses is a very

high percent of clay minerals in the clay- and

silt-sized fractions; montmorillonite predomi-

The clast component of the debris flow is

characterized by an overwhelmingly high percent-

probably vary somewhat in mineralogical composi-

age of volcanic materials. These materials

tion, and are known to vary greatly in size.

angularity, hardness, and color. Many clasts

are blocks and masses 20-25 m in maximum dimen-

sion and a few are as large as 90 m in maximum

dimension. Even 50 km southwest from the cal-

dera, one block 40 m in at least one dimension

airfield. Clasts are angular to subrounded and

rocks probably largely andesite; numerous clasts

andesite, (3) highly fractured red andesite, (4)

yellow to orange-yellow largely clayey masses of

composed commonly of several types of volcanic

have multiple hues. Types of clasts include:

(1) slightly fractured unaltered, hard, dense

variably fractured brown andesite, (5) soft,

hydrothermally altered andesite, (6) highly

fractured green or light-purple andesite, and

(7) pumice. In some localities there are sub-

stantial numbers of boulder-sized clasts of a

diamicton thought to be of glacial origin. No

large concentrations of breadcrusted volcanic

fragments were observed. However, Miller and

Smith (1976) and T. P. Miller and others (writ-

ten commun. 1979), do report bombs in some expo-

sures of volcanic material they term an ash

equivalent of part of the Chetaslina volcanic

debris flow. We have found no wood or charcoal

flow, possibly because vegetation may have been

flow. Although paleomagnetic studies have not

been made on the seemingly autobrecciated volca-

nic masses in the deposit, such studies may be

of value in determining the emplacement tempera-

tures of some of the volcanic clasts (Hoblitt

and Kellogg, 1979). Paleomagnetic studies of

selected volcanic clasts in the deposit along

In some places more than half of the

bles and cobbles of nonvolcanic rock types;

deposit is made up of rounded to subrounded peb-

striated pebbles and cobbles are present in some

Distribution of the deposits

of the Chetaslina debris flow, as determined

ground-surface exposures, and a few test wells

is shown on the map. Below altitudes of about

west of Mount Wrangell and locally along the

ing approximately 12 km northeast-southwest by 70 km northwest-southeast for a total of 840

915 m the remnants are widely distributed south-

lower Chitina River valley over an area measur-

largely from stream-bluff exposures, some

Present known distribution of the remnants

the East Fork Chetaslina River are underway

(K. M. Johnson, oral commun., 1979).

of these exposures.

fragments directly associated with the debris

very sparse on the terrain crossed by the

flow. The ash flow may overlie a lateral

rocks, bombs, and cobble- or larger sized pumice

black andesite, (2) slightly fractured red

was noted 3 km south of the Chitina municipal

pebble size), and the median size of the

ranging from -0.8 to 3.0 phi; the median size of

Chetaslina volcanic debris flow as well as

deposits representing, possibly as many as four This is unfortunate because the conclusion that glacial advances (Nichols, 1965). the debris flow is largely a valley-filling event cannot be more than partly substantiated. General description of deposits A variety of fluvial sediments (chiefly tancolored silty fine sand and some nonvolcanic-The Chetaslina volcanic debris-flow rich pebble gravel) seems to be the most common material underlying the flow near the southdeposits generally are unconsolidated to partly consolidated diamictons having a strong bimodaleasternmost remnants along the Chitina River. In addition to the sand and gravel, a clastity in size of clasts versus size of fragments in the matrix. A matrix is almost totally sparse diamicton is exposed below the debris lacking in some exposures, which gives an flow at a few places along the Kuskulana openwork aspect to the debris-flow deposits. River. Sandy gravel lies beneath the flow at the few exposures of underlying material along The deposits are unsorted, unstratified, and apparently ungraded. the Tonsina River. The matrix component of the debris flow

A wider variety of materials overlies the debris flow. Along much of the middle reach of the Kotsina River, and at some places along the Chetaslina River, a lava flow, part of the Wrangell Lava (Mendenhall, 1905), appears to directly overlie the debris flow (Schrader and Spencer, 1901, p. 59). The lava is described as a generally dark gray to black hypersthene andesite as much as 45 m thick (Mendenhall, 1905, p. 57; H. L. Foster, oral commun., 1965). Locally the lava flow may be of basaltic composition, (Miller and Smith, 1976; T. P. Miller and others, written commun., 1979). At one locality along the north side of the Kotsina River near the mouth of Loraine Creek, an openwork volcanic rubble separates the debris flow from the overlying andesite flow. Volcanic-rich cobble- and boulder-gravel deposits overlie the debris flow at various places along the Chitina River, along the middle reach of the Chetaslina River, and along the Tonsina River near the mouth of Dust Creek. Elsewhere, deposits that overlie the debris flow include (1) sand, (2) diamicton presumably of glacial origin, and (3) sand and gravel composed of mostly nonvolcanic

Thickness of the Chetaslina volcanic

debris-flow deposits below altitudes of about

the average thickness is about 15 m; the

nants are no more than 5-10 m thick.

12.6 km3.

915 m varies greatly. A very rough estimate of

greatest thicknesses are along (1) the southwest

side of the Copper River near the mouth of the

lower reach of the Tonsina River (137 m) and (3)

the middle reach of the Chetaslina River valley

mated. However, considering the areal distribu-

average thickness as 0.015 km, the volume of the

Stratigraphic relations

the Chetaslina debris flow are generally poorly

exposed, especially the underlying materials.

Geologic materials overlying and underlying

tion of the flow remnants as 840 km², and the

debris flow below 915 m altitude may be about

(50 m). At many other locations the flow rem-

The volume of the Chetaslina volcanio

debris flow can only be very roughly esti-

Tonsina River (115 m), (2) the north side of the

The generalized distribution of the andesite flow as shown on the map is based largely on geomorphology and river alinements. In addition to spreading out in the upper valley of the East Fork Chetaslina River and the upland area between the upper Cheshnina River and Long Glacier, the lava probably spread westward beyond Dadina Lake, and possibly spread (1) southwestward to the Copper River and locally to the Tonsina River, and (2) southward along the west side of the Copper River to the Chitina

Speculation on the origin of the Chetaslina volcanic debris flow as a mudflow

Many cycles of volcanic activity have affected the Wrangell Mountains during Quaternary time; in the Mount Drum area, where many K-Ar dates have been determined, two major cycles of activity have been identified as occurring about 525,000 and 325,000 years ago (Richter and others, 1979; T. P. Miller and others, written commun., 1979). The relationship of only a few Quaternary volcanic events to regional glacial events is known. At least three and possibly four glacial events are represented in river bluff exposures along the Copper River and some of its major tributaries (Ferriams and Nichols, 1965; Nichols, 1965). Reworking of deposits representing these glacial events and a mixing of the deposits with products of fluvial and volcanic events has produced a complex geologic record. The record becomes more complex as the distance from volcanic and glacial source areas increases. No single characteristic of the Chetaslina volcanic debris flow is clearly indicative of its origin. However, observations above about 915 m altitude were very reconnaissance in nature, and critical exposures may have been

From a general consideration of the character of the Chetaslina volcanic debris flow and its apparent association with Mount Wrangell, and our examination of geologic literature describing recognition of types of volcanic debris flows (Crandell, 1971; Crandell and Mullineaux, 1976; Sparks, 1976; and references contained therein), it is concluded that the debris flow is best categorized as a volcanic mudflow that locally picked up and incorporated a large variety of other geologic materials. Principal attributes of the debris flow are (1) the widespread distribution, (2) the great size variation and heterogeneity of its volcanic rock

types, (3) the high percentage of soft, hydro-

thermally altered material that probably facili-

tated progressive weakening of the crest of the ancestral volcano, landsliding, and flowage, and

(4) the locally high percentage of nonvolcanic

materials.

Suggested sequence of volcanic and related events involving principally the Chetaslina volcanic debris flow

As an initial consideration, a simple

series of events is tentatively proposed to account for the deposition of the Chetaslina volcanic debris flow and the occurrence of related volcanic activities. However, it is plausible that subsequent work will show that these events are the products of a much more complex and possibly multiple series of volcanic events. It is tentatively suggested that a large phreatic explosion and (or) collapse of the top of ancestral Mount Wrangell may have begun the series of events that initiated the massive, widespread Chetaslina volcanic debris flow (see also Miller and Smith, 1976). Any large-scale explosive and (or) collapse event could have mobilized huge quantities of debris. Hydrogen sulfide gas emanations and clay deposits at existing active small craters along the rim of the caldera (Benson, 1968, p. 146) exemplify the type of hydrothermal activity that probably produced the soft, clayrich altered volcanic rock so widely prevalent in the debris flow. The clayey material would have facilitated flowage of the abundant fragments of rock down the volcano. Heated volcanic debris would have been able to melt ice and snow during its descent from the crest of the volcano (we envision regional glaciers and icecaps similar in size to those of today). The resulting water would further facilitate travel of material down glacier and stream valleys. Another factor that promoted mobility of the debris was the relatively steep gradient of 85 m/km between the caldera and the axis of the Copper River

Blockage of streams by the Chetaslina debris flow may have caused sizable temporary lakes. Volcaniclast-laden cobble and boulder gravel that overlies the flow material near the mouth of the Tebay River and the mouth of Dust Creek on the Tonsina River implies powerful discharge from blocked streams. Likewise, along the west side of the Chetaslina River, the farthest upstream exposure of the debris flow is in part directly overlain by volcanic-rich boulder and cobble gravels. Deposits reflecting blockage of the ancestral Copper River have not been found possibly because of removal by erosion or because the ancestral course of the Copper River was southwest of its current alinement and partly coincident with the present lower course of the Tonsina River.

Possibly the final major event in this proposed sequence of events is the eruption of mostly andesitic lava from several different vents along the caldera rim and possibly from auxiliary vents elsewhere. We believe the lava spread outward from Mount Wrangell into tributaries of the Copper and lower Chitina Rivers since there is no geomorphic evidence of lowlying vents at the base of the mountains or in the adjacent southeastern part of the Copper River basin. In places the lava flow probably extended beyond presently observed outcrops, and subsequently, was eroded differentially by streams and glaciers. Distribution of andesite flows possibly may have determined the present courses of parts of the Tonsina, Dadina, East Fork Chetaslina, and Kotsina Rivers, as well as the Copper River in the area between about Lower Tonsina and Chitina. The alinement of these rivers was probably influenced also by the Chetaslina debris flow. If the lava flow temporarily blocked the Copper River, the flow may have forced the river westward to cut or reoccupy the bedrock channel in which Chitina is

If later volcanic events occurred, they probably resulted in much more localized lava flows, debris flows, or tephra falls. Some examples of young-looking volcanic features are the lava-covered highland south of the glacier which heads the East Fork Chetaslina River (J. R. Williams, oral commun., 1979) and the Mount Zanetti cinder(?) cone northwest of the Mount Wrangell caldera. Both features give the appearance of youthfulness, and might be several thousands of years old or less. The present volcanic activity of Mount Wrangell that continues at several localities near the top of the volcano includes active fumaroles and a few active cinder cones that emit some gases and tephra; a locally high heat flow melts ice and snow in some localities (Benson and Motyka,

Age of the debris flow

The absolute ages of the Chetaslina volcanic debris flow, and volcanic features like the overlying lava flow has not been determined. It is relatively clear, however, that the debris flow did not occur during a major glaciation because most of the deposits observed underlying the debris-flow deposits are indicative of a fluvial environment. It is also apparent that a glacial advance of major proportions differentially eroded the lava flow that overlies the debris flow. Two K-Ar age determinations on plagioclase in the andesite that apparently overlies the debris flow near the mouth of Loraine Creek indicate an age of no older than about 200,000 years at the 95-percent confidence level (J. D. Obradovich, written commun., 1979, U.S. Geological Survey samples DKA 3763 and 3810). Therefore the age of the underlying debris flow is probably only slightly older.

PROPERTY OF DGGS LIBRARY

MISCELLANEOUS FIELD STUDIES MAP MF-1209 CHETASLINA VOLCANIC DEBRIS FLOW, ALASKA

Table 1.—Mineral constituents of matrix of selected Chetaslina volcanic debris-flow deposits. Matrix apparently was hydrothermally altered before being transported

[Identifications by X-ray diffraction; sample 8, identification also by petrographic methods. Values (in estimated parts in ten) give a very general indication of relative amounts of the mineral present. tr, trace; ----, not present or no analysis; <, less than; +, slightly more

d deposited (see map for sample localities)

Constituent minerals	Size fraction and percent of total sample ¹							
	Sample 72			Sample 2c3			Sample 8*	
	sand 43.6	silt 40.2	clay 16.3	sand 44.2	silt 38.1	clay 15.8	clay(?) and silt(?)	
								Plagioclase feldspar
Quartz	8	5	tr	<1	tr			
Mica	1	tr	tr					
Mixed-layer chlorite-montmorillonite						tr?		
Montmorillonite	1	4	5	1+	3+	9	Present.	
Kaolinite	tr	tr	tr				A kaolinitic clay	
							present, possibly dickite.	
Vermiculite			tr					
Clinoptilolite				<1	<1			
Alunite							Present.	
Jarosite		1	4	tr	<1	<1		
Gypsum	tr			<1			Present.	

Particle-size range of fractions: sand, about 0.062-2 mm, -1.0-4.0 phi; silt, 0.002-0.062 mm, 4.0-9.0 phi; and clay, (0.002 mm, > 9.0 phi.

²Sample 7, field no. 60AYe-304A; analyst H. C. Starkey, U.S. Geological Survey Lab. No. DEG-17. 3Sample 2c, field no. 60AYe-158C; analyst H. C. Starkey, U.S. Geological Survey Lab. No. DEG-279. *Sample 8, field no. 54ANi-125; petrographic inspection by R. L. Smith, X-ray analyst Fred Hildebrand, U.S. Geological Survey Lab. No. IWM-764, Job. No. 2236.




Figure 1.--Cumulative size-gradation curves of samples of matrix of selected Chetaslina volcanic debris-flow deposits. No samples were from materials that were



REFERENCES Benson, C. S., 1968, Glaciological studies on Mount Wrangell, Alaska, 1961: Arctic,

v. 21, no. 3, p. 127-152 Benson, C. S., and Motyka, R. J., 1976, Preliminary results of research at the active summit of Mt. Wrangell, Alaska [abs.]: EOS (American Geophysical Union Transactions) v. 57, no. 2, p. 88. 1977. Glacier-volcano interactions--Mt Wrangell, in Annual report 1975-1976: Fairbanks, University of Alaska Geophysical Institute, p. 114-115.

1979, Glacier-volcano interactions on Mt Wrangell, Alaska, in Annual report 1977-1978: Fairbanks, University of Alaska Geophysical Institute, p. 1-25. Bingham, D. K., 1967, Ice motion and heat flow

studies, Mt. Wrangell, Alaska: Fairbanks,

University of Alaska M.S. thesis, 117 p.

Crandell, D. R., 1971, Postglacial lahars from Mount Rainier Volcano, Washington: U.S. Geological Survey Professional Paper 677 Crandell, D. R., and Mullineaux, D. R., 1976, Technique and rationale of volcanic-hazards

appraisals in the Cascade Range, northwestern United States: Environmental Geology, v. 1, no. 1, p. 23-32. Deininger, J. W., 1972, Petrology of the

Wrangell volcanics near Nabesna, Alaska: Fairbanks, University of Alaska M.S. thesis, 66 p.

Dunn, Robert, 1909, Conquering our greatest volcano: Harper's Monthly Magazine, v. 118, p. 497-509, [March]. Ferrians, O. J., Jr., and Nichols, D. R., 1965

Resume of Quaternary geology of the Copper River basin, in Guidebook for Field Conference F, central and south central Alaska: International Association for Quaternary Research Congress, 7th, Lincoln, Nebraska 1965, p. 93-114. Ferrians, O. J., Jr., Nichols, D. R., and

Schmoll, H. R., 1958, Pleistocene volcanio mudflow in the Copper River basin, Alaska [abs.]: Geological Society of America Bulletin, pt. 2, v. 69, no. 12, p. 1563. Furst, F. I., 1968, The reconnaissance petrology of andesites from the Mt. Wrangell caldera, Alaska: Fairbanks, University of Alaska

M.S. thesis, 83 p. Hoblitt, R. P., and Kellogg, K. S., 1979, Emplacement temperatures of unsorted and unstratified deposits of volcanic rock debris as determined by paleomagnetic techniques: Geological Society of America Bulletin, v. 90, pt. 1, no. 7, p. 633-642. Mendenhall, W. C., 1903, The Wrangell Mountains, Alaska: National Geographic Magazine, v. 14, no. 11, p. 395-407.

_____1905, Geology of the central Copper River region, Alaska: U.S. Geological Survey Professional Paper 41, 133 p. Miller, T. P., and Smith, R. L., 1976, Ash flows associated with Wrangell volcano, in The U.S. Geological Survey in Alaska--

Accomplishments during 1975: U.S. Geologi cal Survey Circular 733, p. 52. Nichols, D. R., 1965, Glacial history of the Copper River Basin, Alaska [abs.]: International Association for Quaternary Research, International Congress, 7th Boulder, Colorado, 1965 [Proceedings, v.

Nichols, D. R., and Yehle, L. A., 1969, Engineering geologic map of the southeastern Copper River basin, Alaska: U.S.

Geological Survey Miscellaneous Investigations Map I-529, scale 1:125,000. Richter, D. H., Smith, R. L., Yehle, L. A., and Miller, T. P., 1979, Geologic map of the Gulkana A-2 quadrangle, Alaska: U.S. Geological Survey Geologic Quadrangle Map GQ-1520, scale 1:63,360.

Schrader, F. C., and Spencer, A. C., 1901, The geology and mineral resources of a portion of the Copper River district, Alaska: U.S. Geological Survey, 94 p. [special publica-

of 1941 at Wrightwood, southern California: Geological Society of America Bullein, v. 64, no. 5, p. 547-560. Sparks, R. S. J., 1976, Grain-size variations in

Sharp, R. P., and Nobles, L. H., 1953, Mudflow

ignimbrites and implications for the transport of pyroclastic flows: Sedimentology, v. 23, no. 1, p. 147-188. Wentworth, C. K., 1922, A scale of grade and class terms for clastic sediments: Journa

of Geology, v. 30, no. 5, p. 377-392. Yehle, L. A., 1980, Preliminary surficial geologic map of the Valdez C-1 quadrangle, laska: U.S. Geological Survey Miscellaneous Field Investigations Map MF-1132.

CONTOUR INTERVAL 200 FT (61 M) DATUM IS MEAN SEA LEVEL

INTERIOR-GEOLOGICAL SURVEY, RESTON, VIRGINIA-1980 For sale by Branch of Distribution, U.S. Geological Survey, Box 25286, Federal Center, Denver, CO 80225